

Analytical derivation of lateral pressure profile from microscopic model of lipid bilayer

Sergei I. Mukhin and Svetlana Baoukina

Theoretical Physics Department, Moscow Institute for Steel and Alloys 4 Leninsky pr., 119049
Moscow Russia
e-mail: sergeimoscov@online.ru

Field-theoretical method is proposed, that yields analytic expression for lateral pressure distribution across hydrophobic core of a bilayer lipid membrane. Lipid molecule, repelling entropically surrounding neighbors, is modeled as a flexible string embedded between confining walls in the self-consistent harmonic potential. At room temperature the pressure profile is expressed solely via lowest energy eigenfunctions of the operator of the string energy density. Analytical results compare favorably with molecular dynamics simulations. The theory is suited to study protein-lipid interaction.

PACS numbers:87.16.Dg, 87.14.Cc, 31.15.Qg, 05.20.-y

Development of microscopic theory of fluid lipid membrane is a nontrivial problem. One of the fundamental difficulties lies in the elongated flexible structure of the constituting lipid molecules that results in their large conformational entropy. Neighboring lipids essentially reduces the manifold of accessible conformations of the molecule's hydrocarbon chains in a bilayer membrane as compared with a free lipid molecule. This reduction, in turn, leads to effective entropic repulsion between lipid molecules of non-local nature. Entropic interaction⁴ between hydrocarbon chains causes unusual for common fluids inhomogeneous lateral pressure distribution in thermodynamic equilibrium state. The lateral pressure profile strongly influences the functioning of proteins embedded in the membrane^{2,7}, for instance by shifting their activation barriers^{10,11}. Due to a nanometer scale of membrane thickness the pressure profile is difficult to measure. Several attempts to attack the present problem theoretically have already been made. Some of them involved molecular dynamics simulations of a lipid bilayer^{8,3}. The others exploited numerical methods (e.g. Monte Carlo) with mean-field approach, introducing lateral pressure profile as a substitute for inter-molecular entropic repulsion^{2,5,1}. In the present work we use path-integral technique in combination with adiabatic invariants theorem⁶ and find lateral pressure profile in analytical form. Due to its generality our method permits, in principle, analytical derivation of the lateral pressure profile for a great variety of membrane constituents and lipid variations (head group type, chain length, etc.) by appropriate change of the imposed boundary conditions and membrane spontaneous curvature. Hence, our theory, besides fundamental interest, may be used as a tool for theoretical prediction of the properties of a wide variety of lipid membranes. A sketch of lipid bilayer membrane is given in Fig.1. Hydrocarbon chains form hydrophobic part in lipid molecules. Collisions between chains together with excluded volume effect give rise to entropic repulsion between lipid molecules. This repulsion is responsible for inhomogeneous lateral pressure in the hydrophobic core of the membrane. The ensemble of chains is described within mean-field approximation using microscopic model of a single hydrocarbon chain in effective potential, see 1. The chain is modeled as a flexible string (see Fig. 2) that possesses bending rigidity, K_f . The effective potential, which allows for entropic repulsion between neighboring chains in each monolayer is expressed via lateral pressure distribution $\Pi(z)$ at the depth z multiplied by area $\pi R^2(z)$ swept by the chain in $\{x, y\}$ -plane: $U_{eff} = \pi \Pi R^2(z)$ (compare e.g.¹). Here the chain deviation from the z -axis is $R^2 = R_x^2 + R_y^2$. The energy functional of a single hydrocarbon chain E_t is chosen in the following form:

$$E_t = \int_0^L \left\{ \frac{K_f}{2} \left[\frac{\partial^2 R(z)}{\partial z^2} \right]^2 + \pi \Pi(z) R^2(z) \right\} dz. \quad (1)$$

Here L is chain length; z is coordinate along the chain axis; $\vec{R}$ is vector in $\{x, y\}$ plane characterizing chain deviation from z -axis. In the functional (1) the bending energy term contains second

derivative over z -coordinate rather than over the contour length of the chain. The approximation made is valid provided that chain deviations from the z -axis are small with respect to chain length L (applicability of this condition at room temperature is justified below). We rewrite the energy functional E_t using self-adjoint operator, $\hat{H}$, that is obtained after integrating by parts the expression in (1):

$$E_t = \sum_{i=x,y} \int_0^L R_i(z) \hat{H} R_i(z) dz ; \hat{H} = \frac{1}{2} K_f \frac{\partial^4}{\partial z^4} + \pi \Pi(z). \quad (2)$$

The boundary conditions are: i) chain angle is fixed in the head group region; ii) zero torque is applied at the free chain end; iii) no total force is applied to the chain at head group and at the free chain end. These conditions are expressed as: $R'(0) = R'''(0) = R''(L) = R'''(L) = 0$. An arbitrary chain conformation described by "shape-functions" $R_{x,y}(z)$ can be expanded over orthonormal set of the eigenfunctions, $R_n(z)$, of operator $\hat{H}$:

$$R_x = \sum C_n R_n(z). \quad (3)$$

Since x - and y -deviations are independent we shall retain for simplicity only one index, e.g. x . Then, the energy functional E_t of hydrocarbon chains given in (2) takes equivalent form:

$$E_t = \int_0^L \sum_n E_n C_n^2 R_n^2(z) dz, \quad (4)$$

where E_n are the eigenvalues of the operator $\hat{H}$ in (2).

To find the distribution of lateral pressure along z -axis we shall use the following formal trick⁹. Imagine that we press the membrane laterally at depth z and measure its response. This would give us local lateral pressure. Mathematically this procedure can be performed using the adiabatic invariants theorem⁶. Consider the energy (refenhe) as a sum of potential energies of harmonic oscillators labeled by index n and coordinate z that have rigidities $k_n = 2E_n$ and vibration amplitudes $X_n = C_n R_n(z)$ ¹². Then, the lateral pressure is a sum of individual contributions produced by the oscillators defined above. We calculate each contribution as a thermodynamically averaged force exerted by a one-dimensional oscillator on the surrounding "walls". Assume that the oscillator is centered at $x = 0$ and the walls are positioned at points $\pm d(t)$, where $d(t) = d - vt$ and $v \rightarrow 0$. Here d is z -independent radius of a cylinder, which on average hosts a single chain in a flat bilayer, see Fig.2. The rate of the energy change $d\epsilon_{nz}/dt$ of n -th oscillator at depth z is related to the force, $f_{n,z}$, exerted by the oscillator on the walls: $d\epsilon_{nz}/dt = 2v f_{n,z}$. Alternatively, the rate of the energy change can be calculated using the method of adiabatic invariants⁶:

$$d\epsilon_{nz}/dt = v \frac{\partial I / \partial d}{\partial I / \partial \epsilon_{n,z}} \quad (5)$$

where time-independent adiabatic invariant is $I = \int p dx$. Here $p = \sqrt{2m(\epsilon_{n,z} - k_n x^2/2)}$ is momentum, x is coordinate of the oscillator (mass m of the oscillator does not enter the final expression). Integration in I gives area inside the closed orbit in phase-space $\{p, x\}$ crossing at points $\pm d(t)$ the x -axis and at $\pm p(d(t))$ the p -axis. Equating the above two expressions for $d\epsilon_{nz}/dt$ and taking limit $v \rightarrow 0$ we find the force:

$$f_{n,z} = \begin{cases} \frac{\sqrt{k_n \epsilon_{nz}} \sqrt{1 - d^2 k_n / 2 \epsilon_{nz}}}{\sqrt{2} \arcsin(d \sqrt{k_n / 2 \epsilon_{nz}})}, & d^2 k_n / 2 \epsilon_{nz} < 1, \\ 0, & d^2 k_n / 2 \epsilon_{nz} > 1. \end{cases} \quad (6)$$

The force produced by n -th oscillator at depth z is found as a result of averaging over C_n :

$$\langle f_n(z) \rangle = \int_{-\infty}^{\infty} f_n(\epsilon_{nz}(C_n)) P(C_n) dC_n, \quad (7)$$

where $\epsilon_{nz}(C_n) = E_n C_n^2 R_n^2(z)$ and the distribution function equals:

$$P(C_n) = (E_n/\pi k_B T)^{1/2} \exp \left\{ -\frac{1}{k_B T} \int_0^L E_n C_n^2 R_n^2(z) dz \right\}. \quad (8)$$

Summation over n permits to find the total force at depth z :

$$\langle f(z) \rangle = \sum_n \frac{1}{\sqrt{\pi k_B T R_n^2(z)}} \int_0^\infty \frac{f_n(\epsilon)}{\sqrt{\epsilon}} \exp \left\{ -\frac{\epsilon}{R_n^2(z) k_B T} \right\} d\epsilon, \quad (9)$$

Obtained formula is remarkable: though probabilities for "oscillators" n with different z 's are distributed independently according to formula (8), nevertheless, factors $R_n(z)$ in formula (9) keep track of the fact that "oscillators" are different degrees of freedom of one and the same vibrating string. Actually, only the oscillators that reach the walls at $\pm d$ contribute to the force $f(z)$. This bounds energy ϵ from below: $\epsilon \geq d^2 k_n/2$. Hence, the lower limit of integration in (9) increases fast with index n , because bending part of total energy in (1) results in power-law dependence: $k_n = 2E_n \propto n^4$. Simultaneously, there is an upper limit of integration in (9) $\sim R_n^2(z) k_B T$ imposed by the Boltzmann factor. Hence, non-zero integration interval exists only for the first few terms in (9). Here we limit ourselves to the terms $n = 0, 1$ and find the thermodynamically averaged force:

$$\langle f(z) \rangle \approx \sum_n \frac{k_B T}{2d} R_n^2(z). \quad (10)$$

As the membrane is laterally isotropic, this force can be considered as a radial force in the $\{x - y\}$ plane. Then, the pressure distribution is obtained dividing the radial force in (10) by half the perimeter πd :

$$\Pi(z) \approx \sum_n \frac{k_B T}{2\pi d^2} R_n^2(z). \quad (11)$$

This is the central result of our paper, since, we had shown that the lateral pressure profile that enters the chain energy functional E_t in Eq. (1) is expressed merely via the eigenfunctions of the energy density-operator $\hat{H}$. Using this result we find the corresponding self-consistent solution for $\Pi(z)$ analytically at room temperature. From the estimates for the chain bending rigidity: $K_f \sim k_b T l_p$, where l_p is chain persistence length, it proves that the separation between the eigenvalues E_n of the operator $\hat{H}$ is large with respect to $k_B T$. This justifies approximate projection of the unknown eigenfunctions $R_n(z)$ on the reduced basis formed by the lowest energy eigenfunctions of the free-chain operator $\hat{H}_0 = K_f/2(\partial^2/\partial z)$:

$$R_0 = a_1 Y_0 + b_1 Y_1; \quad R_1 = a_2 Y_0 + b_2 Y_1, \quad (12)$$

where:

$$Y_0(z) = 1/\sqrt{L}, \quad Y_{n \geq 1}(z) = \sqrt{2/L} \{ \cos(\lambda_n z) + \cos(\lambda_n L) \cosh(\lambda_n z) / \cosh(\lambda_n L) \},$$

and the eigenvalues are given by:

$$W_0 = 0, \quad W_{n \geq 1} = K_f / (2L^4) (\pi/4)^4 (4n + 3)^4,$$

where $\lambda_{n \geq 1} L = 3\pi/4 + \pi n$. The coefficients a_i, b_i satisfy the conditions of orthonormality of the functions $R_n(z)$:

$$a_i^2 + b_i^2 = 1, \quad a_1 b_1 + a_2 b_2 = 0.$$

To find the lowest energy eigenfunctions and eigenvalues of the energy-density operator in (2) we substitute unknown functions $R_n(z)$ in the form (12) into expression for $\Pi(z)$ in (11), and insert the resulting expression for $\Pi(z)$ into the equation for eigenfunctions of the operator $\hat{H}$ in (2). We arrive at the following system of non-linear algebraic equations for the lowest eigenvalues E_n of the operator $\hat{H}$ and coefficients a_i, b_i :

$$\begin{cases} a_1\Pi_{00} + b_1\Pi_{01} = a_1E_0 \\ b_1W_1 + b_1\Pi_{11} + a_1\Pi_{10} = b_1E_0 \\ a_2\Pi_{00} + b_2\Pi_{01} = a_2E_1 \\ b_2W_1 + b_2\Pi_{11} + a_2\Pi_{10} = b_2E_1 \end{cases} \quad (13)$$

where the matrix elements Π_{mn} of the "potential energy" operator $\pi\Pi(z)$ are taken on the set of eigenfunctions $Y_n(z)$ defined above. It proves to be that when $\Pi_{00}/W_1 \leq 10$ the system of equations (13) can be solved in closed form. The solutions are:

$$E_0 = \Pi_{00}(1 - I_3\delta), \quad E_1 = W_1 + \Pi_{00}(1 + I_4 + I_3\delta), \quad (14)$$

$$(a_1; b_1) = (1 + \delta^2)^{-1/2} (1; -\delta), \quad (a_2; b_2) = (1 + \delta^2)^{-1/2} (\delta; 1). \quad (15)$$

Here, the following parameters have been introduced:

$$\begin{aligned} I_3 &= L^{3/2} \int_0^L Y_1^3(z) dz \approx -0.4, \quad I_4 = L^2 \int_0^L Y_1^4(z) dz \approx 1.8, \\ \delta &\equiv I_3\Pi_{00}/(W_1 + \Pi_{00}I_4) \approx -0.17. \end{aligned} \quad (16)$$

The matrix element Π_{00} obeys normalization condition:

$$\Pi_{00} = \frac{1}{L} \int_0^L \pi\Pi(z) dz = \frac{\pi P_{eff}}{L}.$$

Effective surface tension in the bilayer, P_{eff} , balances entropic repulsion between hydrocarbon chains. Substituting (15) into (12) and subsequently (12) into (11) we find analytical expression for the lateral pressure profile:

$$\Pi(z) = \frac{P_{eff}}{2(1 + \delta^2)} \{ (Y_0(z) - \delta Y_1(z))^2 + (Y_1(z) + \delta Y_0(z))^2 \}. \quad (17)$$

Normalized lateral pressure profile (17) is plotted in Fig.3. The system parameters are: chain full length $L = 20A$, area per chain in all-trans conformation $A = 20A^2$, chain bending rigidity $K_f = k_B T l_p = 2.8 \cdot 10^{-5}$ erg·cm, $T = 300K$, k_B - Boltzmann constant, $l_p \sim L/3$ - chain persistence length, $P_{eff} \approx 150$ dyne/cm⁸. The calculated profile, shown in Fig.3, qualitatively agrees with the results of the recent molecular dynamics simulations^{8,3}. The behaviour of the $\Pi(z)$ curve can be understood comparing fluctuations of a lipid chain in bilayer and in empty space. In the bilayer the chain fluctuations are strongly suppressed by the neighboring chains with respect to a free lipid molecule. Suppression of chain fluctuation leads to the decrease of conformational entropy and, in turn, results in the enhancement of entropic pressure. The more suppressed the fluctuations, the higher the entropic pressure. The most significant relative decrease of fluctuation amplitude takes place at chain free end ($z \sim L$); at head group region ($z \sim 0$) the relative decrease is noticeable too as lipid heads are not fixed; at mid-chain region the decrease of fluctuations is less pronounced due to segment connectivity in the chain. Now we can validate the exploited approximation of small chain deviations in the bilayer. We calculate the thermodynamic average of chain fluctuation amplitude using expansion (3) and distribution function (8):

$$\langle R^2(z) \rangle = 2 \langle R_{x,y}^2(z) \rangle = k_B T \sum_n \frac{R_n^2(z)}{E_n}. \quad (18)$$

Since, $E_n \propto n^4$, the sum in (18) converges fast. Allowing for relations $R_0^2(z) \sim 1/L$, $E_0 \sim \Pi_{00} \sim P_{eff}/L$, we find rough estimates for the small parameter:

$$\sqrt{\langle R^2(z) \rangle}/L \sim (k_B T/L^2 P_{eff})^{1/2} \sim 0.1. \quad (19)$$

To summarize, we have demonstrated that lateral pressure profile in the hydrophobic core of lipid bilayer is expressed via the eigenfunctions of the density operator of the chain energy. The self-consistent solution for the lateral pressure distribution is found.

The authors acknowledge valuable discussions with Prof. Yu. Chizmadzhev and useful comments at theoretical seminar at Institute Lorentz . The work of S.I.M. was in part supported by RFBR grant 02-02-16354. The work of S.B. was supported by the Foundation "Dynasty".

- ¹ A. Ben-Shaul, in Structure and Dynamics of Membranes, Elsevier Science (1995).
- ² R.S. Cantor, Biophys. J. 76, 2625 (1999a).
- ³ J. Gullingsrud, K. Schulten, Biophys. J. 86, 3496 (2004).
- ⁴ W. Helfrich. Z. Naturforsch. A 33, 305 (1978).
- ⁵ D. Harries, A. Ben-Shaul. J. Chem. Phys. 106 (4), 1609 (1997).
- ⁶ L.D. Landau and E.M. Lifshitz, Mechanics, Pergamon Press, (1960).
- ⁷ D. Marsh, Biochem. Biophys. Acta, 1286, 183 (1996).
- ⁸ E. Lindahl, O. Edholm. J. Chem. Phys. 113 (9), 3882 (2000).
- ⁹ S.I. Mukhin, unpublished.
- ¹⁰ E. Perozo, A. Kloda, D. M. Cortes, and B. Martinac. Nat. Struct. Biol. 9, 696 (2002).
- ¹¹ S.I. Sukharev, W.J. Sigurdson, C. Kung, F. Sachs. J. Gen. Physiol. 113, 525 (1999).
- ¹² A dimensional multiplier of a unit of length is omitted in front of E_n above. Formally, it should be introduced by the transition from the integration over z to the summation. Nevertheless, this multiplier merely leads to the redefinition of the force calculated in (10) as being rather a force per unit of length.




FIG. 1. Mean-field approximation for inter-chain interactions in lipid bilayer.




FIG. 2. Hydrocarbon chain as a flexible string of finite length. $\vec{R}$ is vector in $\{x, y\}$ characterizing deviation of the chain from z-axis; $A = \pi \langle \vec{R}^2 \rangle \equiv \pi d^2$ is average area per string.




FIG. 3. Calculated lateral pressure distribution in the hydrophobic core of the bilayer. The pressure arises from entropic repulsion between hydrocarbon chains in each monolayer. z is coordinate along chain axis spanning from $z = 0$ at head group to $z = L$ at free chain end. Pressure is normalized by $\Pi_0 = P_{eff}/2L$.